Supplementary Table 3. Key KAP indicators disaggregated by health facility (%)

Indicator (variable; criterion)	Guay.	Perq.	Panch.	Pto.Lib.	S.J.Non.	Total
KNOWLEDGE						
Adequate PNC knowledge (C11; ≥2)	55.0	55.0	60.0	54.0	63.9	57.6
Early initiation (C12; =1)	100.0	96.7	100.0	100.0	100.0	99.3
Adequate test knowledge (C16; ≥2)	16.7	33.3	25.0	28.6	21.3	25.0
Fe/FA knowledge partial+ (C19; ≥1)	98.3	98.3	100.0	98.4	98.4	98.7
Safe delivery place (C20; =1)	98.3	100.0	98.3	98.4	98.4	98.7
Adequate contracep. knowledge (C27; ≥2)	20.0	45.0	25.0	28.6	44.3	32.6
ATTITUDES (% agree+strongly agree)						
Medical visits important (C28)	100.0	100.0	100.0	100.0	100.0	100.0
Mother-baby health monitoring (C29)	100.0	100.0	100.0	100.0	100.0	100.0
PNC before 3rd month (C31)	100.0	100.0	100.0	98.4	100.0	99.7
Blood tests (C32)	98.3	100.0	100.0	100.0	98.4	99.3
Blood pressure (C33)	100.0	100.0	100.0	100.0	100.0	100.0
Ultrasound (C34)	100.0	98.3	96.7	100.0	100.0	99.0
Nutrition recommendations (C35)	100.0	98.3	96.7	100.0	98.4	98.7
Fe/FA good for mother-baby (C36)	100.0	100.0	100.0	100.0	100.0	100.0
Rejects home delivery (C38)	91.7	100.0	95.0	96.8	95.1	95.7
Contraception important (C42)	88.3	98.3	95.0	100.0	91.8	94.7
PRACTICES						
Regular PNC visits (C44; =1)	98.3	91.7	98.3	95.2	98.4	96.4
Prior contraceptive use (C56; =1)	58.3	63.3	73.3	79.4	54.1	65.8
Vaccination ≥2 doses (C47)	35.0	11.7	41.7	39.7	36.1	32.9
Fe/FA never forgets (C37; =1)	76.7	55.0	60.0	55.6	70.5	63.5

Notes: Percentages by UCSF. Variable and calculation criterion shown in parentheses. Guay.=Guaymango (n=60); Perq.=Perquín (n=60); Panch.=Panchimalco (n=60); Pto.Lib.=Puerto de La Libertad (n=63); S.J.Non.=San Juan Nonualco (n=61). Attitude items were measured using a four-point Likert scale coded as 1=strongly agree, 2=agree, 4=disagree and 5=strongly disagree. A neutral midpoint was included during pilot testing and removed before the final survey; therefore, response code 3 does not appear in the dataset.